

Effect of mixed edible coatings containing gum tragacanth and Aloe vera on postharvest quality of strawberries during storage

Aryou Emamifar^{1*}, Sudabeh Bavaisi²

Received: 2017.02.02

Accepted: 2017.08.07

Abstract

The effects of mixed coating based on aloe vera (AG) and gum tragacanth (GT) on the microbial, physicochemical and sensorial properties of fresh strawberries were evaluated during 20 days of storage (1 °C, 95 % RH) compared to uncoated fruits. The coating solutions were prepared by mixing solution of AG diluted 1:3 with distilled water and GT solution (0.6 % w/v in distilled water) at different concentrations (25 % AG +75 % GT, 50 % AG +50 % GT and 75 % AG +25 % GT). Microbial stability (fungi (yeasts and molds) and total aerobic bacteria), physicochemical characteristics (ascorbic acid (AA), weight loss, firmness, titratable acidity, soluble solid content (SSC), anthocyanin content, total phenolic and antioxidant activity) and sensory attributes (color, taste, odor and overall) of the samples were evaluated after 0, 4, 8, 12, 16 and 20 days of storage compared to uncoated fruits. Comparing with untreated fruits, 50 % AG +50 % GT treatment significantly ($p < 0.05$) decreased the microbial growth, weight loss and AA degradation and maintained firmness, anthocyanin and phenol contents and antioxidant activity. A greater visual acceptance was observed in fruits treated with 50 % AG + 50 % GT. The combination of AG/GT solution as a proper coating formulation in addition to have high antimicrobial activity; have great potential to extend shelf life of fresh strawberries.

Keywords: Strawberry; Aloe vera; Gum tragacanth; Shelf life.

Introduction

Strawberry fruits have a short shelf life, so after harvest, the marketability of them will reach about 5 d at temperature between 0 and 4 °C with 90-95 % relative humidity (Jiang *et al.* 2001). The loss of fresh strawberries can reach 40 % from harvesting to consumption (Park *et al.* 2005). There have been many studies on reducing postharvest losses of strawberry fruits using pre or post-harvest treatments including: polyamines (Ponappa *et al.* 1993), methyl jasmonate (Ayala-Zavala *et al.* 2005), UV (Erkan *et al.* 2008), nitric oxide (Wills *et al.* 2000), 1-methylcyclopropene (Bower *et al.* 2003) ultrasound (Aday *et al.* 2013), salicylic acid (Zhang *et al.* 2010) as well as application of edible coating (Tanada-Palmu and Grosso 2005; Del-Vallea *et al.*

2005). Application of edible coating on perishable fruits such as strawberries or fresh cut fruits is one of the most common methods to increase their shelf life through reducing the respiration rate, water loss, improving the sensory attributes and retarding the microbial growth (Bifani *et al.* 2007). Alginate, cellulose, chitosan, chitin, lipids, mucilage, milk protein, starch, wax, and zein have been widely used as an edible coating in the food industry (Valverde *et al.* 2005). Gum tragacanth is one of the most popular herbal gum that exudates from stem of *Astragalus gummifer*. Safety assessment and approving their use in foods (emulsifying agent, stabilizer, filing and thickening agent) have been confirmed (E-number E413) by the Scientific Committee for Food of the European Community (Gavlighi *et al.* 2013; Samanta *et al.* 2010; Fayazzadeh *et al.* 2014). Gum tragacanth is an acidic hetero-polysaccharide that contained some protein-bounded polysaccharides (Saffari *et al.* 2013). It consists of two different fractions including tragacanthin (water-soluble) and bassorin (water insoluble) with different rheological

1. Assistant Professor, Department of Food Science and Technology, Buali-Sina University, Hamedan, Iran

2. Master in Horticultural Science, Department of Horticultural Science, University of Kurdistan, Sanandaj, Iran.

(*Corresponding author email: a.emamifar@basu.ac.ir)

DOI: 10.22067/iftstrj.v1396i3.62317

properties (Balaghi *et al.* 2010). Aloe vera (*Aloe barbadensis* Miller) is a natural plant with functional properties such as antibacterial, antifungal and antioxidant activity (Nejatzadeh-Barandozi 2013). The transparent gel of aloe vera that contains nutritional components has also edible and medicinal values (Zafari *et al.* 2015). Two main components obtained from the fresh harvested aloe vera leaves including liquid fraction containing bitter excaudate and semisolid fraction containing parenchyma tissue. Moreover, it has improved postharvest life quality of fruits such as grape, nectarine, papaya and pomegranate arils (Valverde *et al.* 2005; Navarro *et al.* 2011). Many researches have been done on the applications of gum tragacanth in food formulations. Recently, it has been used in food packaging (Mostafavi *et al.* 2016). There are few reports on the use of tragacanth as an edible coating. The main mechanism of the film formation in polysaccharide films is the breaking apart polymer segments and reforming of the polymer chain into film matrix or gel that achieved by several film-forming mechanisms including solvent evaporation, creating hydrophilic and hydrogen binding or electrolytic and ionic crosslinking (Park *et al.* 2014). The safety of biopolymer materials, solutions and additive that used as fruit coatings is very important for its application on the food industry (Šuput *et al.* 2015). The main purpose of this research is to evaluate the effects of gum tragacanth combined with aloe vera gel on the microbial contamination, physicochemical and sensory properties of fresh strawberries during cold storage.

Materials and Methods

In the first step, the optimal concentration of diluted gum tragacanth in distilled water as an edible coating solution for fresh strawberries was determined and in the next step, the effects of the mixed edible coating containing aloe vera (AG) and gum tragacanth (GT) at different GT/ AG ratio on the postharvest quality of strawberries were evaluated.

Plant material

Under the guidance of expert botanists, at the University of Kurdistan, fresh strawberries (*Fragaria X anannasa* Duch.), cv. 'Parous', were harvested at maturity stage (based on 80 % red color on the surface) from a commercial farm in Sanandaj, Iran. They were immediately pre-cooled (at 1°C for 1 hour) after harvest and were directly transported to the laboratory. Then, fresh fruits were sorted for uniform size, color, and removed unripe and damaged and then were stored at 1 °C and 95 % RH.

Preparation of coating formulation (Gum tragacanth solution)

The ribbon tragacanth gum (slight-yellowish clear appearance) was purchased from the local market in Sanandaj, Iran. They were grinded (National, K039131, Cixi City, China) and sieved (Retch sieve, 0.425 mm, Lincoln, U.K.) to produce fine powder. The powdered gum tragacanth at different concentrations (0, 0.3, 0.6 and 0.9% w/v; of total solid) was added gently to the distilled water, mixed for 1 h and let stand overnight at ambient temperature to fully hydrate. Gum tragacanth solutions were pasteurized in a water bath at 70°C for 45 min and were cooled to 25°C (Marpudi *et al.* 2011).

Coating and storage

Strawberry fruits were immersed and the coating treatments were performed at 25°C by immersing the strawberries for 5 min in gum tragacanth coating solutions and dried on a sterile stainless-steel screen under fans for about 1 h. Gum tragacanth solutions were pasteurized in a water bath at 70°C for 45 min and were cooled to 25°C (Marpudi *et al.* 2011). Sterile distilled water was used as the control solution. The coated and uncoated (Control) fruits were then packaged in polyethylene packages (72 packages, each containing 15 pieces of strawberries) and stored at 1°C with 95 % RH. The samples were evaluated at three replications (Each replicate consisted of one packed with 15 plants) for their microbiological, physicochemical, and sensory characteristics

at 0, 4, 8, 12, 16 and 20 d of storage. After the selection of the best concentration of gum tragacanth solution, it was applied for preparing the mixed coating solution with aloe vera gel solution.

Preparation of coating formulation (Aloe vera gel)

Aloe vera leaves were cut from greenhouse-grown plants (University of Kurdistan, Sanadaj, Iran). They were washed with water, disinfected (in a 2% sodium hypochlorite solution for 30 min) and rinsed with distilled water. Aloe vera gel matrix was then separated from the outer cortex of leave, ground, and filtered to produce a fresh aloe vera gel. After pasteurization (at 70°C for 45 min) the gel was cooled to 25°C (Marpudi *et al.* 2011). The coating solution was prepared by aloe vera gel diluted 1:3 with sterile distilled water (Valverde *et al.* 2005; Hassanpour 2015).

Preparation of coating formulation (Aloe vera and Gum tragacanth solutions)

Mixtures of pasteurized (GT) and aloe vera (AG) solutions at different concentrations (25% AG+ 75 % GT, 50% AG+ 50% GT and 75% AG+ 25% GT) were prepared as an edible coating. Fruits were dipped in coating solutions or sterile distilled water as the control solution, at 25°C for 5 min. They dried on a sterile stainless-steel screen under fans for about 1 h to ensure surface dryness. The coated and uncoated (control) fruits were packaged (72 packages, each containing 15 pieces of strawberries) and stored (under the same conditions as mentioned above). Samples were evaluated exactly according to section 2.3. (Coating and storage).

Microbiological evaluations

At each sampling time, for each replicate a mixed sample with quarter of each fruit was transferred aseptically to a stomacher bag and homogenized. Decimal dilutions of the suspension of strawberries in sterile peptone water (0.1%) were prepared. With the spread plate and pour plate methods, enumeration of total aerobic bacteria (at 30°C for 2 d), yeasts and molds (at 25°C for 2d) were carried out on the plate count agar (PCA, Scharlau Chemie,

S.A., Barcelona, Spain) and the potato dextrose agar (PDA, Scharlau Chemie, S.A., Barcelona, Spain), respectively. Serial dilutions (10^{-2} to 10^{-3}) were performed in triplicate and dilutions were plated in duplicate. The results were expressed as colony-forming units (CFU) per gram (Sogvar *et al.* 2016).

Physicochemical properties

Weight loss (WL) and Firmness

The weight of each packaged (replicate) was recorded immediately after packaging and after each 4d of storage. Weight loss was expressed as the percentage loss of the initial fresh weight. The firmness of strawberries in each box replicate was evaluated by texture analyzer (Santam, STM-1) with an 8 mm probe with constant speed of 20 mm min⁻¹. Two different measurements were done on opposite sides of the central zone of each fruit (Sogvar *et al.*, 2016). Force values were expressed as newton (N).

Titrateable acidity (TA), soluble solid content (SSC) and Ascorbic acid

A mixed sample of all fruit with one-eighth segments of each fruit per replicate juiced together and used for measuring the titrateable acidity (TA), soluble solid content (SSC) and ascorbic acid. TA was measured based on titrimetric method by using 0.1 M NaOH (to pH 8.1) and was expressed as percentage of citric acid per 100 mL fruit juice (AOAC, 2002). SSC was detected by digital refractometer (AOAC, 2002). Ascorbic acid was determined based on titrimetric method by 2,6-dichlorophenolindophenol as a titrant (AOAC 967.21) and expressed as mg kg⁻¹ fresh weight basis of fruits (AOAC, 2002).

Total anthocyanin content (TAC), total phenolic concentrations (TPC) and Total antioxidant activity (TAA)

At the sampling time, a bulked sample of all fruits with 1/8 th of each fruit per replicate was cut, frozen and stored at -80°C and they were used for determination of total anthocyanin content (TAC), total phenolic

(TP) concentrations and total antioxidant activity. Total anthocyanin content was measured by a pH differential method (Cheng and Breen 1991). The absorbance of the combined sample extract with quarter of each fruit was read at 510 nm and 700 nm in buffers at pH 1.0 and 4.5. Results were expressed as milligrams of pelargonidin 3-glucoside (P 3-G) equivalents per kilogram of fresh weight. Total phenolic concentrations were determined according to Folin Ciocalteu method using gallic acid as the standard curve and the results were presented as milligram of gallic acid equivalent per kilogram of fruit fresh weight (Singleton *et al.*, 1999). Measurement of the total antioxidant activity of frozen samples was carried out by 2, 2-diphenyl-1-picrylhydrazil (DPPH) radical-scavenging method (Sanchez Moreno *et al.* 1999). The absorbance was measured at 517 nm, using a spectrophotometer (UNICO UV-2100, Shanghai, China) and the results presented as the percentage inhibition of DPPH radicals.

Sensory analysis

Sensory characteristics (color, odor, taste, and overall) of uncoated and coated strawberries were evaluated based on a 5-point scale (extremely like= 5, moderately like= 4, neither liked nor disliked= 3, moderately dislike= 2 and extremely dislike= 1) by 10 expert panelists. Fruits were sampled at 0, 4, 8, 12, 16 and 20 days of storage (Emamifar *et al.* 2010).

Statistical analysis

Analysis of variance was carried out using the SAS statistical software release 6.12 (SAS Institute, Cray, NC) based on completely randomized designs. Significant differences among the data were represented as $p < 0.05$.

Results and Discussion

Experimental section 1

Table 1, shows the variations of the mean microbial populations, weight loss and ascorbic acid content in the tragacanth coated fruits compared to control (uncoated) during storage (at 1°C and 95% RH). In all samples,

the microbial populations and weight loss increase and ascorbic acid content decreases at different rates as the storage time increases, significantly ($p < 0.05$). The mean initial populations of yeasts and molds and total aerobic bacteria in freshly harvested strawberries immediately after packaging (control samples) were found to be 1.47 log CFU g⁻¹ and 1.38 log CFU g⁻¹, respectively. For the coated strawberry fruits with 0.6% gum tragacanth solution, significance decreases were observed in yeasts and molds and total aerobic bacteria populations compared to all other coated and uncoated samples, after 20 d. The weight loss of uncoated fruits was increased significantly ($p < 0.05$) up to 34.15% as compared to 26.58, 18.85 and 22.38 % for strawberries coated with 0.3, 0.6 and 0.9% gum tragacanth solutions, respectively. Ascorbic acid degradation in control samples was higher than in other coated samples after 20 d of storage. At the end of storage period, strawberries coated with 0.6% gum tragacanth solution maintained higher ascorbic acid content (64%) compared to other that coated with 0.3% gum tragacanth solution (52%), 0.9% gum tragacanth solution (58%) and control (46%), significantly ($p < 0.05$). However, the 0.6% w/v solution of gum tragacanth treatment shows a significant ($p < 0.05$) effect on reducing the microbial populations and weight loss and maintaining the ascorbic acid content in strawberry fruits as compared to the other treatments and control during storage. These results are in agreement with the findings of Mohebbi *et al.* (2012). Treviño-Garza *et al.* (2015) found that edible active coatings based on polysaccharides reduced the microbial growth in coated strawberry fruits and increased their shelf life up to 15 d. Gol *et al.* (2013) showed that the combination of Hydroxypropyl methyl cellulose (HPMC) and chitosan as an edible coating can maintain the ascorbic acid content in fresh strawberries during cold storage. Significant reductions of weight loss were observed in strawberries coated with cassava starch (Garcia *et al.* 2012) and chitosan-based edible coatings (Han *et al.*

2004) during cold storage. Strawberry fruits coated with 0.6% w/v gum tragacanth solution showed significantly ($p < 0.05$) higher values of firmness, total anthocyanin, total phenol,

antioxidant activity and sensory attributes and lower values of SSC and TA compared to other treatments and control (data not shown).

Table 1. Effect of tragacanth (mean \pm sd) ^a on the yeasts and molds, total aerobic bacteria populations, weight loss and ascorbic acid of strawberries stored at 1 °C for 20 d.

Tragacanth concentration (%)	Storage time(days)						
	0	4	8	12	16	20	
Molds and Yeasts (log CFU g ⁻¹)	0	1.47 ^k \pm 0.05	2.25 ^g \pm 0.04	2.61 ^e \pm 0.01	2.81 ^{cd} \pm 0.04	3.04 ^b \pm 0.02	3.37 ^a \pm 0.05
	0.3	1.47 ^k \pm 0.04	1.88 ⁱ \pm 0.03	2.29 ^g \pm 0.05	2.57 ^e \pm 0.05	2.78 ^d \pm 0.02	2.89 ^c \pm 0.07
	0.6	1.47 ^k \pm 0.04	1.61 ^j \pm 0.03	1.93 ⁱ \pm 0.02	2.08 ^h \pm 0.05	2.39 ^f \pm 0.05	2.57 ^e \pm 0.02
	0.9	1.47 ^k \pm 0.01	1.62 ^j \pm 0.07	2.06 ^h \pm 0.01	2.39 ^f \pm 0.05	2.61 ^e \pm 0.07	2.74 ^d \pm 0.02
Total aerobic bacteria (log CFU g ⁻¹)	0	1.38 ^m \pm 0.03	2.13 ^{gh} \pm 0.01	2.45 ^d \pm 0.04	2.70 ^c \pm 0.01	2.91 ^b \pm 0.03	3.29 ^a \pm 0.05
	0.3	1.38 ^m \pm 0.07	1.84 ^{ij} \pm 0.03	2.04 ^h \pm 0.03	2.26 ^{ef} \pm 0.06	2.54 ^d \pm 0.01	2.78 ^c \pm 0.08
	0.6	1.38 ^m \pm 0.06	1.54 ^l \pm 0.04	1.76 ^j \pm 0.02	1.91 ⁱ \pm 0.05	2.19 ^{fg} \pm 0.08	2.48 ^d \pm 0.07
	0.9	1.38 ^m \pm 0.02	1.65 ^k \pm 0.06	1.91 ⁱ \pm 0.01	2.16 ^g \pm 0.04	2.34 ^e \pm 0.02	2.70 ^c \pm 0.02
Weight loss (%)	0	0.00 ^j \pm 0.00	6.75 ⁱ \pm 0.70	15.76 ^e \pm 0.36	21.10 ^c \pm 0.40	26.64 ^b \pm 0.45	34.15 ^a \pm 1.01
	0.3	0.00 ^j \pm 0.00	6.10 ⁱ \pm 0.28	12.90 ^f \pm 0.65	18.09 ^d \pm 0.60	21.57 ^c \pm 0.22	26.58 ^b \pm 0.17
	0.6	0.00 ^j \pm 0.00	5.20 ⁱ \pm 0.69	9.12 ^h \pm 0.44	12.51 ^{fg} \pm 0.48	15.69 ^e \pm 0.18	18.85 ^d \pm 0.19
	0.9	0.00 ^j \pm 0.00	5.95 ^k \pm 0.09	10.85 ^{gh} \pm 0.13	15.04 ^e \pm 0.34	17.78 ^d \pm 0.24	22.38 ^c \pm 0.22
Ascorbic acid (mg kg ⁻¹)	0	555.50 a \pm 3.34	435.50 e \pm 11.1	382.25 fg \pm 7.79	347.75 ji \pm 1.12	307.75 l \pm 3.36	255.50 m \pm 11.12
	0.3	555.50 a \pm 5.57	487.70 c \pm 8.86	425.50 e \pm 2.22	371.15 gh \pm 1.16	335.50 jk \pm 1.19	294.40 i \pm 3.39
	0.6	555.50 a \pm 1.12	532.25 b \pm 2.23	488.87 c \pm 3.34	422.25 e \pm 7.72	385.50 fg \pm 8.88	355.50 hi \pm 1.16
	0.9	555.50 a \pm 7.78	592.20 b \pm 1.14	455.50 d \pm 3.32	395.50 f \pm 6.63	355.51 hi \pm 3.34	326.65 k \pm 1.09

Values followed by the same letter in the same row are not significantly different ($p < 0.05$).

Experimental section 2

Microbiological evaluation

Strawberry fruits can become contaminated with spoilage microorganisms during growth in the field, harvesting and postharvest handling, or during storage (Barth *et al.* 2009). Moreover, the fruit surface has long been considered a suitable environment for the growth of microorganisms. More generally, fruits surface microbes have impact on the rates of food spoilage (Akhtar *et al.* 2016). Freshly harvested strawberries contain more nutrients and high water content. Therefore, they are highly susceptible to microbial decay. There are multiple factors that considerably affect microbial decay during storage including temperature, relative humidity and O₂ concentration (Kader 1998). Edible coating act as barriers to moisture and oxygen during storage and delayed decay development in coated fruits as compared to

uncoated (Valverde *et al.* 2005). The variations of the mean microbial populations in coated fruits compared to control (uncoated) during cold storage are shown in Fig. 1. For all samples, the microbial population increased significantly ($p < 0.05$) at different rates as the storage time increased. The populations of yeasts and molds and total aerobic bacteria in control samples reached 3.81 log CFUg⁻¹ and 3.82 log CFUg⁻¹, respectively, after 20 d. According to Fig. 1. The mixed gum based treatments have significant ($p < 0.05$) effects on retarding microbial growth in all strawberry samples during cold storage. In comparing all treated and untreated samples, 50 % GT + 50 % AG treatment showed the most antifungal (2.72 log CFU g⁻¹) and antibacterial (2.78 log CFU g⁻¹) activity on fresh strawberries during 20 d of storage at 1 °C. The effect of the volume ratio of AG to GT on antimicrobial activity of mixed coating material is

pronounced significantly ($p < 0.05$). As shown in Fig. 1, increased AG: GT ratio is believed to be related to the decreased antimicrobial activity of coating materials. However, coating formulation containing lower concentrations of aloe vera (50%) have significantly more antimicrobial activity than coating formulation containing 75% of aloe vera ($p < 0.05$). Aloe vera contains several special compounds with antimicrobial activity, including anthraquinones (Garcia-Sosa *et al.* 2006), dihydroxy- anthraquinones (Wu *et al.* 2006) and saponins (Reynolds and Dweck 1999). Ferro *et al.* (2003) showed that aloe vera gel exhibited an antimicrobial activity against gram positive and gram-negative bacteria as well as yeasts and molds. Moreover, the antifungal activity of the aloe vera gel against fungi spores such as *Penicillium*, *Botrytis* and *Alternaria* and against mycelium growth of fungi such as *Rhizoctonia*, *Fusarium* and *Colleotrichum* (Saks and Barkai-Golan 1995; Jasso de Rodriguez *et al.* 2005) have reported.

Shelf life extension and microbial load reduction of pomegranate arils, table grape and sweet cherry using aloe vera gel as an antimicrobial coating have been reported (Martínez-Romero *et al.* 2013; Valverde *et al.* 2005; Martínez-Romero *et al.* 2006). Recently, antimicrobial activity of gum tragacanth against some gram negative and gram-positive bacteria and molds and yeasts has been reported (Ghayempour *et al.* 2015; Singh *et al.* 2015). Muthulakshmi (2013) showed that the interaction between gum tragacanth amino groups with gram-negative bacterial cell wall as well as the binding of gum tragacanth carboxylate groups with the gram-positive bacterial cell and the fungal cell wall are the most important reasons for antimicrobial activity of this gum. Moreover, the complex structure and high molecular weight of gum tragacanth have more positive effects on antimicrobial activity of gum tragacanth (Muthulakshmi 2013).




Fig. 1. Effect of mixed coating based on aloe vera (AG) and gum tragacanth (GT) on the yeasts and molds populations (a) and total aerobic bacteria populations (b) of strawberries stored at 1 °C for 20 d. Vertical bars represent standard error ($n = 3$).

Weight loss

According to Fig. 2.A., in all samples, the weight loss increases at different rates as the storage time increases, significantly ($p < 0.05$). Strawberries coated with 50% GT+ 50% AG showed the lower rate of weight loss compared to the others, so that after 20 d of storage, the weight loss of uncoated fruits reached to

26.6% as compared to 21.1% (25% AG+ 75% GT), 18.7 % (50% AG+ 50% GT) and 19.3% (75% AG+ 25% GT), respectively (Fig. 2.A.). No significant differences were observed in weight loss between two-treatment coating including 50% AG+ 50% GT and 75% AG+ 25% GT, except on the 8th d ($p < 0.05$). Weight loss often occurred due to water evaporation in

fruits and vegetables during respiration process (Yaman and Bayoindirli 2002). However, the main impact of coatings application are based on their hygroscopic properties, which form a water barrier layer between fruits and environments and reduce the fruit surface evaporation (Mohebbi *et al.* 2012). Aloe vera coating has also been reported to reduce the weight loss of table grapes, sweet cheery, apple and papaya stored at cold storage (Valverde *et al.* 2005; Martinez-Romero *et al.* 2006; Marpudi *et al.* 2011; Ergun and Satici 2012). Mohebbi *et al.* (2012) reported that the combination of aloe vera and gum tragacanth are more effective on reducing the weight loss of button mushroom. It can be concluded that the combination of aloe vera and gum tragacanth in the specified concentration ranges have a synergetic effects on reducing the weight loss of the strawberries during cold storage.

Firmness

The firmness is one of the most important physical quality characteristics of fruit during storage (Pasquariello *et al.* 2013). As shown in Fig. 2.B., with increasing the storage time the rate of softening increased in all of strawberries, but it was faster in uncoated than those of coated samples (Fig. 2.B.). After 20 d, firmness value decreased from 3.05 N to 1.41, 1.73, 2.26 and 1.93 N in uncoated and coating treatments including 25% AG+ 75% GT, 50% AG+ 50% GT and 75% AG+ 25% GT, respectively (Fig. 2.B.). 50% AG+ 50% GT coating was more effective in firmness retention of the strawberries compared to others ($p < 0.05$). These results are in agreement with the results reported by many studies which stated that the firmness of strawberry fruits treated with edible coating including cactus mucilage (Del-Valle *et al.* 2005), starch (García *et al.* 2012) and gluten (Tanada-Palmu and Grosso 2005) were maintained during storage. Degradation of the middle lamella of the cortical parenchyma cells and cell separation and subsequent microbial infection are the main reasons for the softening of strawberry fruits during storage (Rahman *et al.*

2016).

Titration acidity (TA) and soluble solids content (SSC)

For all samples, SSC increased and TA decreased during storage (Fig. 2.C and D). The variations of TA and SSC values in uncoated fruits during storage were significantly ($p < 0.05$) higher than those of coated fruits. According to Fig. 2C. and D, after 20 d of storage; the highest TA (0.705%) and the lowest SSC (6.9%) were recorded in strawberries treated with 50% AG+ 50% GT, compared to the other treatments and control samples, indicating uncoated strawberries presented a more pronounced ripening development than coated strawberries, similarly to that found in starch-coated strawberry (Mali and Grossmann, 2003). These results agreed with those found by EL Gaouth *et al.* (1991), Pelayo *et al.* (2002) and Koyuncu (2004). Zheng *et al.* (2007) showed that decreasing in titratable acidity of strawberry fruits during cold storage caused by increasing their respiration rate. The increase in SSC of strawberry fruits during storage could be related to degradation of cell-wall polymers or other polysaccharides such as hemicelluloses and water loss (Hernandez-Munoz *et al.* 2008; Tanada-Palmu and Grosso 2005).

Ascorbic acid content

Ascorbic acid content of strawberries can be influenced by pre-harvest and postharvest factors including cultivar, growing season, environmental conditions, harvest time and storage conditions (Skrovankova *et al.* 2015; Cordenunsi *et al.* 2005). The average ascorbic acid content of strawberry fruits is ranged from 190 mg kg⁻¹ to 715 mg kg⁻¹ (Lee and Kader 2000). According to Fig. 3.A., ascorbic acid in all coated and uncoated samples reduced over 20 d of storage. Coating containing 50% AG+ 50% GT was the most effective treatment to maintain ascorbic acid (29.26%) in strawberries during storage up to 20 d while this value was decreased to 17.29 % for uncoated and 23.31 and 28.35% for coating containing 75% AG+ 25% GT and

25% AG+ 75% GT, respectively. Autoxidation is one of the main reasons for ascorbic acid depletion during storage of fruits (Plaza *et al.* 2006). Coating materials provide a protective thin layer between the fresh fruit and its surrounding atmosphere that decrease the

moisture transfer, O₂, and CO₂ exchange (Tapia *et al.* 2008; Atress *et al.* 2010). Adetunji *et al.* (2012) have found that pineapple coated with aloe vera gel showed the shelf life extension by decreasing the oxygen permeability.




Fig. 2. Effect of mixed coating based on aloe vera (AG) and gum tragacanth (GT) on the weight loss (A), Firmness (B), Soluble solid content (C) and Titratable acidity (D) of strawberries stored at 1°C for 20 d. Vertical bars represent standard error (n = 3).

Total anthocyanin content (TAC), total phenolic concentrations (TPC)

Anthocyanin and phenolic compounds are beneficial phytochemicals that affect the functional properties of strawberry fruits (Giampieri *et al.* 2012). There is an increase in consumers' demand of fresh strawberry fruits due to beneficial properties (Flores-Félix *et al.* 2015). It is therefore critical to consider their changes during storage. Coating treatments

and storage time significantly ($p < 0.05$) affected the initial total anthocyanin content (50.5 mg kg⁻¹ of pelargonidin-3-glucoside) and total phenol content (355 mg kg⁻¹ of gallic acid on fresh weight basis) of fresh strawberries during storage (Fig. 3.B. and C.). After 4 d of storage, a sharp increase in anthocyanin was observed in all coated fruits compared to uncoated (Fig. 3. B.). Fruits treated with 50%

AG+ 50% GT had the greatest anthocyanin content than those of other treatments and untreated fruits after 20 d of storage ($p < 0.05$). The amount of anthocyanin content of strawberry is dependent on the cultivar, storage temperature and O_2 concentration (Cordenunsi *et al.* 2005). Several studies have shown that the biosynthesis of the strawberry fruits have continued during storage (Kalt *et al.* 1999; Cordenunsi *et al.* 2005; Ayala-Zavala *et al.* 2005). However, a slight increase in anthocyanin content for all samples was observed up to 20 d of storage (Fig. 3.B). According to Fig 3.C., the total phenolic contents in all uncoated samples decreased following storage for 20 d (223 mg kg^{-1}) but they decreased in coated samples up to 16 d in 25% AG+ 75% GT (252 mg kg^{-1}), 50% AG+ 50% GT (278.5 mg kg^{-1}) and 75% AG+ 25% GT (271 mg kg^{-1}) and then increased up to 20 d. The total phenolic contents decrease at different rates as the storage time increases, significantly ($p < 0.05$). However, phenol degradation in control was higher than other coated samples after 20 d (Fig.3.C.). Strawberries coated with 50% AG+ 50% GT maintained higher total phenols (295 mg kg^{-1}) compared with other coated treatments at the end of storage. Coating materials stabilize the TSS/ TA ratio in fruits and cause the pH remain at a low level. It is leading to a decrease in enzyme activity of polyphenol oxidase and increase total phenol concentrations (Singha *et al.* 2009). Gol *et al.* (2013) showed that total phenolic contents increased in strawberries during cold storage.

Antioxidant activity

Antioxidant activity in coated and uncoated strawberry fruits decreased during cold storage at 1°C (Fig.3.D.). Reducing the rate of antioxidant activity in coated fruits was faster than that of uncoated. The high antioxidant activity was observed in strawberry fruits that were coated with 25% AG+ 75% GT (45.5%), 50% AG+ 50% GT (55.5%) and 75% AG+ 25% GT (50.6%) compared to control (36.5%) after 20 d. As shown in Fig.3.D, coating containing 50% AG+ 50% GT is the most

effective treatment to maintain antioxidant activity in strawberry fruits during cold storage. However, aloe vera gel has inherent antioxidant capacity resulted in aloe gel coated fruits retained their antioxidant activity. Previous studies have shown that antioxidant capacity of aloe vera at different stages of development, is due to many active compounds with different degrees of antioxidant capacity (Vieira *et al.* 2016; Wu *et al.* 2006). Ascorbic acid and the total phenolic contents in strawberry fruits could have a significant impact on the antioxidant activity (Kelebek *et al.* 2009; Wang and Lewers 2007). Studies have shown that grapes (Serrano *et al.* 2006) and raspberry (Hassanpour 2015) coated with aloe vera gel had higher antioxidant capacity than uncoated.

Sensory evaluation

According to Fig.4. Significant differences ($p < 0.05$) were determined in the sensory characteristics of coated strawberries compared to uncoated except for the color score. Scores for all sensorial attributes fall down as the storage time increased, significantly ($p < 0.05$). After 20 d, the panelists assigned the highest color, odor, taste and overall scores to strawberry fruits coated with 50% AG+ 50% GT, 75% AG+ 25% GT and 25% AG+ 75% GT, respectively and the lowest score being associated with uncoated samples (Fig. 4.). The odor and texture scores correlated well with the microbial load and firmness values presented in Fig. 1. and Fig. 2.B, respectively. It can be concluded that, 50% GT+ 50% AG treatment as the best coating combination, could keep the sensory attributes of strawberries over the storage period. Garcia *et al.* (2012) reported that sensory characteristics of the strawberries coated with cassava starch had been accepted by consumers up to 12 d compared to uncoated fruits. Benitez *et al.* (2013) showed that the coating of kiwifruit slices with 5% aloe vera gel improved their marketability up to 8 d.

Conclusion

The results obtained in the present research showed that application of combined edible coating containing aloe vera and gum tragacanth is a new preservation approach for fresh strawberry fruits at 1°C. Using solution coating containing 50 % diluted aloe vera (1:3) and 50% gum tragacanth solution (0.6 % w/v), extended the shelf life of fresh strawberries up to 20 d without any side

effects on their physicochemical characteristics and sensorial attributes. However, application of combined coating containing aloe vera and gum tragacanth for storage of fresh strawberries is not sufficient for long time storage. This study revealed that for extension shelf life of fresh strawberries, it is necessary to apply another treatment as a new hurdle in addition to coating treatments.




Fig.3. Effect of mixed coating based on aloe vera (AG) and gum tragacanth (GT) on Ascorbic acid (A), Anthocyanin (B), Total phenols (C) and Antioxidant activity (D) of strawberries stored at 1 °C for 20 d. Vertical bars represent standard error (n = 3).




FIG. 4. Effect of mixed coating based on aloe vera (AG) and gum tragacanth (GT) on the sensory attributes of strawberries after 20 d at 1 °C.

References

- Aday MS, Temizkan R, Büyükcan MB, Caner C. (2013). An Innovative Technique for Extending Shelf Life of Strawberry: Ultrasound. *LWT - Food Science Technology* 52, 93–101.
- Adetunji CO, Fawole OB, Arowora KA, Nwaubani SI, Ajayi ES, Oloke JK, Majolagbe OM, Ogundele BA, Aina JA, Adetunji J B. (2012). Effects of Edible Coatings from Aloe Vera Gel on Quality and Postharvest Physiology of *Ananas Comosus* (L.) Fruit during Ambient Storage. *Global Journal of Science Frontier Research Bio-Tech & Genetics* 12, 39-43.
- Akhtar T, et al. (2016). Identification of Microbial Contamination of Popular Fruits of Bangladesh and Assessment the Effects of Alternative Preservatives Instead of Formalin. *American Journal of Microbiological Research* 4 138-142.
- Atress ASH, El-Mogy MM, Aboul-Anean HE, Alsanious BW. (2010). Improving Strawberry Fruit Storability by Edible Coating as a Carrier of Thymol or Calcium Chloride. *Journal of Horticultural Science & Ornamental Plants* 2, 88-97.
- AOAC. (2002). Acidity (Titratable) of Fruit Products. Gaithersburg, MD: Official Methods of Analysis of AOAC International, *Official Method* 942.15.
- AOAC. (2002). Solids (Soluble) in Fruits and Fruit Product: Refractometer Method. Gaithersburg, MD: Official Methods of Analysis of AOAC International, *Official Method* 932.12.
- AOAC. (2002). Vitamin C (ascorbic acid) in Vitamin Preparations and Juices: 2, 6 Dichloroindophenol Titrimetric Method Final Action. Gaithersburg, MD: Official Methods of Analysis of AOAC International, 2002. *Official Method* 967.21.
- Ayala-Zavala JF, Wang SY, Wang CY, Gonzalez-Aguilar GA. (2005). Methyl Jasmonate in Conjunction with Ethanol Treatments Increased Antioxidant Capacity, Aroma Compounds and Postharvest Life of Strawberry Fruit. *European Food Research and Technology* 22, 1438-1443.
- Balaghi S, Mohammadifar MA, Zargaraan A. (2010). Physicochemical and Rheological Characterization of Gum Tragacanth Exudates from Six Species of Iranian Astragalus. *Food Biophysics* 5:1, 59-71.

- Barth M, Hankinson TR, Zhuang H, Breidt F. (2009). Microbiological Spoilage of Fruits and Vegetables. In W.H. Sperber, M.P. Doyle (Eds.), *Compendium of the Microbiological Spoilage* (pp. 257-275). Springer-Verlag New York.
- Benitez S, Achaerandio I, Sepulcre F, Pujola M. (2013). Aloe Vera Based Edible Coatings Improve the Quality of Minimally Processed 'Hayward' Kiwifruit. *Postharvest Biology and Technology* 81, 29–36.
- Bifani V, Ramírez C, Ihl M, Rubilar M, García A, Zaritzky N. (2007). Effects of Murta (*Ugni Molinae Turcz*) Extract on Gas and Water Vapor Permeability of Carboxymethylcellulose-Based Edible Films. *LWT - Food Science and Technology* 40, 1473-1481.
- Bower JH, Biasi WV, Mitcham EJ. (2003). Effect of Ethylene and 1-MCP on the Quality and Storage Life of Strawberries. *Postharvest Biology and Technology* 28, 471-423.
- Cheng G, Breen P. (1991). Activity of Phenylalanine Ammonia-Lyase (PAL) and Concentrations of Anthocyanins and Phenolics in Developing Strawberry Fruit. *Journal of the American Society for Horticultural Science* 116, 865-869.
- Cordenunsi BR, Genovese MI, Nascimiento JRO, Hassimotto NMA, Santos, RJ, Lajolo, FM. (2005). Effects of Temperature on the Chemical Composition and Antioxidant Capacity of Three Strawberry Cultivars. *Food Chemistry* 91: 113– 121.
- Del-Vallea V, Hernández-Muñozb P, Guardac A, Galottod MJ. (2005). Development of a Cactus-Mucilage Edible Coating (*Opuntia Ficus Indica*) and Its Application to Extend Strawberry (*Fragaria Ananassa*) Shelf-Life. *Food Chemistry* 91, 751–756.
- El Gaouth A, Arul J, Ponnampalam R, Boulet M. (1991). Chitosan Coating Effect on Storability and Quality of Fresh Strawberries. *Journal of Food Science* 12, 1618–1632.
- Emamifar A, Kadivar M, Shahedi M, Soleimani-Zad S. (2010). Evaluation of Nanocomposite Packaging Containing Ag and Zn on Shelf Life of Fresh Orange Juice. *Innovative Food Science & Emerging Technologies* 11, 742–748.
- Ergun M, Satici F. (2012). Use of Aloe Vera Gel as Biopreservative for 'Granny Smith' and 'Red Chief' Apples. *The Journal of Animal & Plant Sciences* 22, 363-368.
- Erkan M, Wang YS, Wang YC. (2008). Effect of UV Treatment on Antioxidant Capacity, Antioxidant Enzyme Activity and Decay in Strawberry Fruit. *Postharvest Biology and Technology* 48, 163-171.
- Fayazzadeh E, Rahimpour S, Ahmadi SM, Farzampour S, Anvari MS, Boroumand MA, Ahmadi SH. (2014). Acceleration of Skin Wound Healing with Tragacanth (*Astragalus*) Preparation: An Experimental Pilot Study In Rats. *Acta Medica Iranica* 52,3-8.
- Ferro VA, Bradbury F, Cameron P, Shakir E, Rahman SR, Stimson WH. (2003). In Vitro Susceptibilities of *Shigella Flexneri* and *Streptococcus Pyogenes* to Inner Gel of *Aloe Barbadosis Miller*. *Antimicrobial Agents and Chemotherapy* 43, 1137-1139.
- Flores-Félix JD, Silva LR, Rivera LP, Marcos-García M, García-Fraile P, Martínez-Molina E, Rivas R. (2015). Plants Probiotics as a Tool to Produce Highly Functional Fruits: The Case Of Phyllobacterium And Vitamin C In Strawberries. *PLoS ONE*, 10:4, 1-10.
- García LC, Pereira LM, de Luca Sarantópoulos CIG, Hubinger MD. (2012). Effect of Antimicrobial Starch Edible Coating on Shelf-Life of Fresh Strawberries. *Packaging Technology and Science* 25, 413–425.
- García-Sosa K, Villarreal-Alvarez N, Lubben P, Peña-Rodríguez LM. (2006). Chrysophanol, an Antimicrobial Anthraquinone From The Root Extract of *Colubrina Greggii*. *Journal of the Mexican Chemical Society* 50, 76-78.
- Gavlighi HA, Meyer AS, Zaidel DN, Mohammadifar MA, Mikkelsen JD. (2013). Stabilization of Emulsions By Gum Tragacanth (*Astragalus Spp.*) Correlates to the Galacturonic Acid Content and Methoxylation Degree of The Gum. *Food Hydrocolloids* 31:1, 5-14.
- Ghayempour S, Montazer M, Mahmoudi Rad M. (2015). Tragacanth Gum as a Natural Polymeric

- Wall for Producing Antimicrobial Nanocapsules Loaded With Plant Extract. *International Journal of Biological Macromolecules* 81, 514-520.
- Giampieri F, Tulipani S, Alvarez-Suarez JM, Quiles JL, Mezzetti B, Battino M. (2012). The Strawberry: Composition, Nutritional Quality, and Impact on Human Health. *Nutrition* 28:1, 9-19.
- Gol NB, Patel PR, Rao TR. (2013). Improvement of Quality and Shelf Life of Strawberries with Edible Coatings Enriched with Chitosan. *Postharvest Biology and Technology* 85,185-195.
- Han C, Zhao Y, Leonard SW, Traber MG. (2004). Edible Coatings to Improve Storability and Enhance Nutritional Value of Fresh and frozen Strawberries (*Fragaria X Ananassa*) and Raspberries (*Rubus ideaus*). *Postharvest Biology and Technology* 33, 67-78.
- Hassanpour H. (2015). Effect of Aloe Vera Gel Coating on Antioxidant Capacity, Antioxidant Enzyme Activities and Decay in Raspberry Fruit. *LWT - Food Science and Technology* 60, 495-501.
- Hernández-Muñoz P, Almenar E, Del Valle V, Velez D, Gavara R. (2008). Effect of Chitosan Coating Combined with Postharvest Calcium Treatment on Strawberry (*Fragaria Ananassa*) Quality During Refrigerated Storage. *Food Chemistry* 110, 428-435
- Jasso de Rodriguez D, Hernandez-Castillo R, Rodriguez-Gracia I, Angulo-Sanchez JL. (2005). Antifungal Activity in Vitro of Aloe Vera Pulp and Liquid Fraction Against Plant Pathogenic Fungi. *Industrial Crops and Products* 21, 81-87.
- Jiang YM, Joyce DC, Terry LA. (2001). 1-Methylcyclopropene Treatment Affects Strawberry Fruit Decay. *Postharvest Biology and Technology* 23, 227-232.
- Kalt W, Forney CF, Martin A, Prior RL. (1999). Antioxidant Capacity, Vitamin C, Phenolics and Anthocyanins after Fresh Storage of Small Fruits. *Journal of Agricultural and Food Chemistry* 47:11, 4638-4644.
- Kader AA. (1998). Influence of Preharvest and Postharvest Environment on Nutritional Composition of Fruits and Vegetable. In Quebedeaux B, Bliss FA. (Eds.). *Horticulture and Human Health, Contribution of Fruits and Vegetables*. Prentice-Hall, Englewood Cliffs, N.J.
- Kelebek H, Selli S, Canbas A, Cabaroglu T. (2009). HPLC Determination of Organic Acids, Sugars ,Phenolic Compositions and Antioxidant Capacity of Orange Juice and Orange Wine Made From a Turkish cv.Kozan. *Microchemical Journal* 91:2, 187-192.
- Koyuncu MA. (2004). Quality Changes of Three Strawberry Cultivars during the Cold Storage. *European Journal of Horticultural Science* 65, 193-200.
- Lee SK, Kader AA. (2000). Pre-harvest and Post-Harvest Factors Influencing Vitamin C Content of Horticultural Crops. *Postharvest Biology and Technology* 20:3, 207-220.
- Mali S, Grossmann MVE. (2003). Effects of Yam Starch Films on Storability and Quality of Fresh Strawberries (*Fragaria ananassa*). *Journal of Agricultural and Food Chemistry* 51, 7005-7011.
- Marpudi SL, Abirami LSS, Pushkala R, Srividya N. (2011). Enhancement of Storage Life and Quality Maintenance of Papaya Fruits Using Aloe Vera Based Antimicrobial Coating. *Indian Journal of Biotechnology* 10, 83-89.
- Martínez-Romero D, Castillo S, Guillén F, Díaz-Mul HM, Zapata PJ, Valero D, Serrano M. (2013). Aloe Vera Gel Coating Maintains Quality and Safety of Ready-To-Eat Pomegranate Arils. *Postharvest Biology and Technology* 86, 107-112.
- Martinez-Romero D, Albuquerque N, Valverde, JM, Guillén F, Castillo S, Valero D, Serrano M. (2006). Postharvest Sweet Cherry Quality and Safety Maintenance by Aloe Vera Treatment: A New Edible Coating. *Postharvest Biology and Technology* 39: 93-100.
- Mohebbi M, Ansarifard E, Hasanpour N, Amiroufsefi MR. (2012). Suitability of Aloe Vera and Gum Tragacanth as Edible Coatings for Extending the Shelf Life of Button Mushroom. *Food Bioprocess Technology* 5, 3193-3202.
- Mostafavi RS, Kadkhodae R, Emadzadeh B, Koocheki A. (2016). Preparation and

- Characterization of Tragacanth–Locust Bean Gum edible Blend Films. *Carbohydrate Polymer* 139, 20-27.
- Muthulakshmi S. (2013). Investigation on the Biopolymers Papain, Gum Acacia, Gum Tragacanth and Gum Guar; Physical and Antimicrobial Properties. Thesis submitted on manonmaniam Sundaranar University in partial fulfillment of the requirements for the award of the degree of doctor of philosophy in physics. Reg no 2335.
- Navarro D, Díaz-Mula HM, Guillén F, Zapata PJ, Castillo S, Serrano M, Valero D, Martínez-Romero D. (2011). Reduction of Nectarine Decay Caused by *Rhizopus stolonifer*, *Botrytis cinerea* and *Penicillium digitatum* with Aloe Vera Gel Alone or with the Addition of Thymol. *International Journal of Food Microbiology* 151, 241–246.
- Nejatzadeh-Barandozi, F. (2013) Antibacterial Activities and Antioxidant Capacity of Aloe Vera. *Organic and Medicinal Chemistry Letters* 3, 2-8.
- Park HJ, Byun YJ, Kim YT, Whiteside WS, Bae HJ. (2014). Processes and Applications for Edible Coating and Film Materials from Agropolymers. In J.H. Han (Eds.), *Innovations in Food Packaging*, Second Edition. (pp. 257-275). Elsevier Ltd.
- Park SI, Stan SD, Daeschel MA, Zhao Y. (2005). Antifungal Coatings on Fresh Strawberries (*Fragaria × Ananassa*) to Control Mold Growth during Cold Storage. *Journal of Food Science* 70, 202-207.
- Pasquariello MS, Rega P, Migliozi T, Capuano LR, Scortichini M, Petriccione M. (2013). Effect of Cold Storage and Shelf Life on Physiological and Quality Traits of Early Ripening Pear Cultivars. *Scientia Horticulturae* 162, 341–350.
- Pelayo C, Ebeler SE, Kader AA. (2003). Postharvest Life and Flavor Quality of Three Strawberry Cultivars Kept At 5°C In Air or Air+20kpa CO₂. *Postharvest Biology and Technology* 27, 171-183.
- Plaza L, Sánchez-Moreno C, Elez-Martínez P, De Ancos B, Martín-Belloso O, Cano, M P. (2006). Effect of Refrigerated Storage on Vitamin C and Antioxidant Activity of Orange Juice Processed by High-Pressure or Pulsed Electric Fields with Regard to Low Pasteurization. *European Food Research and Technology* 223, 487–493.
- Ponappa T, Scheerens JC, Miller AR. (1993). Vacuum Infiltration of Polyamines Increases Firmness of Strawberry Slices under Various Storage Conditions. *Journal of Food Science* 58, 361-364.
- Rahman MM, Moniruzzaman M, Ahmad MR, Sarker BC, Alam MK. (2016). Maturity Stages Affect the Postharvest Quality and Shelf Life of Fruits of Strawberry Genotypes Growing in Subtropical Regions. *Journal of the Saudi Society of Agricultural Sciences* 15, 28–37.
- Reynolds T, Dweck AC. (1999). Aloe vera leaf gel: a review update. *J Ethnopharmacol* 68: 3-37.
- Saks Y, Barkai-Golan R. 1995. Aloe Vera Gel Activity Against Plant Pathogenic Fungi. *Postharvest Biology and Technology* 6, 159-165.
- Saffari MM, Farzi M, Emam-Djomeh Z, Moini S, Mohammadifar MA. (2013). Applying Iranian Gum Tragacanth to Improve Textural Properties of Maltodextrin Microcapsules. *Journal of Texture Studies* 44, 12–20.
- Samanta A, Ojha D, Mukherjee B. (2010). Stability Analysis of Primary Emulsion Using a New Emulsifying Agent Gum Odina. *Natural Science* 2:5, 494-505.
- Sanchez-Moreno C, Larrauri JA, Saura-Calixto F. (1999). A Procedure to Measure the Antiradical Efficiency of Polyphenols. *Journal of the Science of Food and Agriculture* 76, 270–276.
- Serrano M, Valverde JM, Guillen F, Castillo S, Martínez-Romero D, Valero D. (2006). Use of Aloe Vera Gel Coating Preserves the Functional Properties of Table Grapes. *Journal of Agricultural and Food Chemistry* 54:11, 3882-3886.
- Singh BR, Dubey S, Siddiqui MZ. (2015). Antimicrobial Activity of Natural Edible Gums. *World Journal of Pharmaceutical Sciences* 3:11, 2217-2221.

- Singha SP, Singha Z, Swinny EE. (2009). Postharvest Nitric Oxide Fumigation Delays Fruit Ripening and Alleviates Chilling Injury during Cold Storage of Japanese Plums (*Prunus salicina Lindell*). *Postharvest Biology and Technology* 53, 101–108.
- Singleton VL, Orthofer R, Lamuela-Raventos RM. (1999). Analysis of Total Phenols and Other Oxidation Substrates and Antioxidants by Means of Folin–Ciocalteu, Reagent. *Methods in Enzymology* 299, 152–178
- Skrovankova S, Sumczynski D, Mlcek J, Jurikova T, Sochor J. (2015). Bioactive Compounds and Antioxidant Activity in Different Types of Berries. *International Journal of Molecular Sciences* 16, 24673–24706.
- Sogvar OB, Koushesh Saba M, Emamifar A. (2016). Aloe Vera and Ascorbic Acid Coatings Maintain Postharvest Quality and Reduce Microbial Load of Strawberry Fruit. *Postharvest Biology and Technology* 114, 29–35.
- Šuput DZ, Lazić VL, Popović SZ, Hromiš NM. (2015). Edible Films and Coatings – Sources, Properties and Application. *Food and Feed Research* 42, 11–22.
- Tanada-Palmu SP, Grosso CRF. (2005). Effect of Edible Wheat Gluten-Based Films and Coatings on Refrigerated Strawberry (*Fragaria. ananassa*) Quality. *Postharvest Biology and Technology* 36, 199–208.
- Tapia MS, Rojas-Graü MA, Carmona A, Rodríguez FJ, Soliva-Fortuny R, Martín- Belloso O. (2008). Use of Alginate- and Gellan-Based Coatings for Improving Barrier, Texture and Nutritional Properties of Fresh-Cut Papaya. *Food Hydrocolloids* 22, 1493–1503.
- Treviño-Garza MZ, García S, del Socorro Flores-González Ma, Arévalo-Niño, K. (2015). Edible Active Coatings Based on Pectin, Pullulan, and Chitosan Increase Quality and Shelf Life of Strawberries (*Fragaria ananassa*). *Journal of Food Science* 80, M1823–M1830.
- Valverde JM, Valero D, Martínez-Romero D, Guillen F, Castillo S, Serrano M. (2005). Novel Coating Based on Aloe Vera Gel to Maintain Table Grape Quality and Safety. *Journal of Agricultural and Food Chemistry* 53, 7807–7813.
- Vieira J, López M, Rodríguez Diana Jasso de, Sousa Maria C, Vicente A, Martins Joana A. (2016). Effect of Chitosan Aloe Vera Coating on Postharvest Quality of Blueberry (*Vaccinium corymbosum*) Fruit. *Postharvest Biology and Technology* 116, 88–97.
- Wang SY, Lewers KS. (2007). Antioxidant Capacity and Flavonoid Content in Wild Strawberries. *Journal of the American Society for Horticultural Science* 132:5, 629–637.
- Wills RBH, Ku VVV, Leshem YY. (2000). Fumigation with Nitric Oxide to Extend the Postharvest Life Of Strawberries. *Postharvest Biology and Technology* 18, 75– 79.
- Wu YW, Ouyang J, Xiao, XH, Gao, WY, Liu Y. (2006). Antimicrobial Properties and Toxicity of Anthraquinones by Microcalorimetric Bioassay. *Chinese Journal of Chemistry* 24, 45–50.
- Yaman Ö, Bayoindirli L. (2002). Effects of An Edible Coating and Cold Storage on Shelf-Life and Quality of Cherries. *LWT-Food Science and Technology* 35, 146–150.
- Yang FM, Li HM, Li F, Xin ZH, Zhao LY, Zheng YH, Hu QH. (2010). Effect of Nano-Packing on Preservation Quality of Fresh Strawberry (*Fragaria ananassa Duch. cv Fengxiang*) During Storage at 4 °C. *Journal of Food Science* 75, 236–240.
- Zafari E, Mohammadkhani A, Roohi V, Fadaei A, Zafari H. (2015). Effect of Exogenous Putrescine and Aloe Vera Gel Coating on Post-Harvest Life of Strawberry (*Fragaria ananassa Duch.*) Fruit, Cultivar Kamarosa. *International Journal of Agriculture and Crop Sciences* 8, 785–791.
- Zhang H, Ma L, Turner M, Xu, H, Zheng X, Dong Y, Jiang, S. (2010). Salicylic Acid Enhances Bio Control Efficacy of *Rhodotorula glutinis* Against Postharvest *Rhizopus* Rot of Strawberries and the Possible Mechanisms Involved. *Food Chemistry* 122, 577–583.
- Zheng Y, Wang SY, Wang CY, Zheng W. (2007). Changes in Strawberry Phenolics, Anthocyanins, and Antioxidant Capacity in Response to High Oxygen Treatments. *LWT-Food Science and Technology* 40:1, 49–57.

تاثیر پوشش خوراکی ترکیبی بر پایه صمغ کتیرا و آلوئه‌ورا بر کیفیت پس از برداشت توت‌فرنگی طی انبارداری

آریو امامی‌فر^{1*} - سودابه باویسی²

تاریخ دریافت: 1395/11/14

تاریخ پذیرش: 1396/05/16

چکیده

اثربودن پوشش ترکیبی بر پایه ژل آلوئه‌ورا و ژل کتیرا بر ویژگی‌های میکروبی، فیزیکوشیمیایی و حسی توت‌فرنگی تازه طی 20 روز انبارداری (دمای یک درجه سانتی‌گراد و رطوبت نسبی 95 درصد) در مقایسه با نمونه بدون پوشش ارزیابی گردید. پوشش‌ها با مخلوط کردن محلول ژل آلوئه‌ورا رقیق شده (به نسبت 1:3 وزنی حجمی در آب مقطر) و ژل کتیرا (با غلظت 0/6 درصد وزنی حجمی در آب مقطر) در غلظت‌های مختلف (25% ژل آلوئه‌ورا + 75% ژل کتیرا، 50% ژل آلوئه‌ورا + 50% ژل کتیرا و 75% ژل آلوئه‌ورا + 25% ژل کتیرا) تهیه شدند. پایداری میکروبی (تعداد کپک و مخمر و کل باکتری‌های مزوفیل هوازی)، خصوصیات فیزیکوشیمیایی (اسید آسکوربیک، کاهش وزن، سفتی، اسیدیته، مواد جامد محلول، محتوی آنتوسیانین، فنل کل و فعالیت ضد اکسایشی) و ویژگی‌های حسی (رنگ، طعم، بو و پذیرش کلی) نمونه‌ها پس از 0، 4، 8، 12، 16 و 20 روز از شروع انبارداری در مقایسه با نمونه بدون پوشش (شاهد) ارزیابی گردید. پوشش‌های حاوی 50% ژل آلوئه‌ورا + 50% ژل کتیرا در مقایسه با نمونه‌های بدون پوشش به صورت معنی‌داری ($p < 0/05$) رشد میکروبی، کاهش وزن و تخریب اسید آسکوربیک را در نمونه‌های توت‌فرنگی کاهش داده و سفتی یافت، محتوی آنتوسیانین، فنل کل و ظرفیت ضد اکسایشی آن‌ها را حفظ کردند. همچنین بیشترین امتیاز ویژگی‌های حسی به توت‌فرنگی‌های پوشش داده با 50% ژل آلوئه‌ورا + 50% ژل کتیرا اختصاص یافت. به هر ترتیب محلول ترکیبی از دو ژل کتیرا و آلوئه‌ورا به عنوان یک فرمول پوششی مطلوب، علاوه بر خاصیت ضد میکروبی زیاد، از توانایی بالایی در افزایش ماندگاری توت‌فرنگی تازه برخوردار بود.

واژه‌های کلیدی: توت‌فرنگی، آلوئه‌ورا، صمغ کتیرا، عمر نگهداری

1- استادیار، گروه صنایع غذایی دانشگاه بوعلی سینا، همدان.

2- دانشجوی کارشناسی‌ارشد، گروه صنایع غذایی دانشکده کشاورزی دانشگاه کردستان.

* - نویسنده مسئول : (Email: a.emamifar@basu.ac.ir)